

# **Endocytic pathways mediating oligomeric A $\beta$ 42 neurotoxicity**

Chunjiang Yu, Evelyn Heath, Kevin Laxton, and Mary Jo LaDu<sup>§</sup>

Department of Anatomy and Cell Biology, University of Illinois at Chicago, Chicago,  
IL 60612

<sup>§</sup>Corresponding author

Email addresses:

CY: [cjyu@uic.edu](mailto:cjyu@uic.edu)

EN: [enwabu1@uic.edu](mailto:enwabu1@uic.edu)

KL: [klaxto2@uic.edu](mailto:klaxto2@uic.edu)

MJL: [mladu@uic.edu](mailto:mladu@uic.edu)

# **Abstract**

## **Background**

One pathological hallmark of Alzheimer's disease (AD) is amyloid plaques, composed primarily of amyloid- $\beta$  peptide ( $A\beta$ ). Over-production or diminished clearance of the 42 amino acid form of  $A\beta$  ( $A\beta_{42}$ ) in the brain leads to accumulation of soluble  $A\beta$  and plaque formation. Soluble oligomeric  $A\beta$  ( $oA\beta$ ) has recently emerged to be as a likely proximal cause of AD.

## **Results**

Here we demonstrate that endocytosis is critical in mediating  $oA\beta_{42}$ -induced neurotoxicity and intraneuronal accumulation of  $A\beta$ . Inhibition of clathrin function either with a pharmacological inhibitor, knock-down of clathrin heavy chain expression, or expression of the dominant-negative mutant of clathrin-assembly protein AP180 did not block  $oA\beta_{42}$ -induced neurotoxicity or intraneuronal accumulation of  $A\beta$ . However, inhibition of dynamin and RhoA by expression of dominant negative mutants, reduced neurotoxicity and intraneuronal  $A\beta$  accumulation. Pharmacologic inhibition of the dynamin-mediated endocytic pathway by genistein also reduced neurotoxicity.

## **Conclusions**

These data suggest that dynamin-mediated and RhoA-regulated endocytosis are integral steps for oligomeric  $A\beta_{42}$ -induced neurotoxicity and intraneuronal  $A\beta$  accumulation.

## Background

Amyloid- $\beta$  peptide ( $A\beta$ ) is believed to be a causative agent underlying the pathological mechanism for Alzheimer's disease, the major form of dementia in the elderly [1]. The levels of soluble  $A\beta$  species appears to correlate with disease progression [2-12]. Evidence points to soluble oligomeric  $A\beta$  (o $A\beta$ ) as the assembly form of the peptide that is likely the proximal cause in AD [13-24], leading to synaptic dysfunction and eventual neuron loss in the vulnerable regions of AD brains (for recent review [25]). Extracellular o $A\beta$  has been proposed to bind the cell surface and several putative receptors have been proposed [26-29], including  $\alpha 7$  nicotinic acetylcholine receptor ( $\alpha 7nAChR$ ) and cellular prion protein. The binding of o $A\beta$  to these receptors may lead to the disturbance of intracellular  $Ca^{++}$  homeostasis, disrupting functions of NMDAR [30, 31] and AMPAR [32, 33], and activating caspases [34].

In addition to extracellular  $A\beta$ ,  $A\beta$  accumulates inside neurons. Intraneuronal  $A\beta$  accumulation has been identified in Down syndrome and AD patients, amyloid precursor protein (APP) and PS1 Presenilin 1 transgenic mice, and cultured cells [35-52]. In AD patients, intraneuronal  $A\beta_{42}$  accumulation appears in vulnerable brain regions prior to extracellular amyloid formation and accumulates with aging [35-41, 43, 48, 49, 53-56]. In addition, synaptic dysfunction occurs prior to, or in the absence of, amyloid plaques in both AD and APP transgenic mouse brains [9, 57-60]. Studies using triple transgenic mice demonstrated that intraneuronal  $A\beta$  causes the onset of early AD-related cognitive deficits [47, 61, 62]. Intriguingly, clearance of intraneuronal  $A\beta$  by immunotherapy rescued early cognitive deficits, prior to changes

in plaque density. Intraneuronal A $\beta$  and cognitive deficits re-emerged with the subsequent withdrawal of immunotherapy [62, 63]. These observations support the hypothesis that intraneuronal A $\beta$  accumulation may be one of the initial steps in a cascade of events leading to AD [64, 65]. Intraneuronal A $\beta$  could be viewed as compromised clearance of extracellular soluble A $\beta$  by neurons. Excessive accumulation of intraneuronal A $\beta$  could lead to cellular organelle dysfunction and eventual neuron death. For example, intraneuronal A $\beta$  was reported to activate caspase 6 leading to neuronal apoptosis [66]. We recently demonstrated that intracellular oA $\beta$ 42 can activate casein kinase-2, causing inhibition of fast axonal transport [67].

Neurons, like many other cell types, have several major endocytic pathways, including clathrin-dependent, caveolae-dependent, and clathrin- and caveolar-independent pathways. However, the specific endocytic pathways involved in oA $\beta$ -uptake and neurotoxicity remain unclear. Using complementary approaches of pharmacological inhibition, genetic manipulation by over-expressing dominant-negative mutants and gene knock-down, we provide data that show that the endocytosis of oA $\beta$ 42 is linked to neurotoxicity via a dynamin-dependent and RhoA-mediated endocytic pathway *in vitro*.

We previously established a homogenous preparation of oA $\beta$ 42 [19] that causes neurotoxicity in co-cultures of primary neurons and glia, as well as Neuro-2A cells (N2A) [18, 68, 69]. This oA $\beta$ 42 preparation also inhibits LTP [70], causes cognitive deficits [71], disrupts fast axonal transport [67], and induces neuroinflammation [72]. Here we focus mainly on endocytic pathways in relation to oA $\beta$ 42 toxicity in N2A cells.

## Results

### **Clathrin-dependent endocytic pathway is not involved in oA $\beta$ 42 mediated toxicity**

As accumulation of intracellular A $\beta$ 42 accompanies neurotoxicity, we wanted to determine whether blocking specific endocytic pathways would inhibit neurotoxicity. We used several approaches to examine clathrin-mediated endocytosis, a major endocytic pathway (refer to Figure 4). First, to directly target clathrin, we transiently transfected N2A cells with siRNA specifically targeting mouse clathrin heavy chain. Western blot analysis showed that siRNA substantially reduced clathrin protein levels, in comparison to non-target siRNA control (Figure 1A inset). However, clathrin siRNA failed to block oA $\beta$ 42 toxicity, similar to the non-targeting siRNA control (Figure 1A). Second, we expressed a dominant-negative mutant of the neuron-specific clathrin-assembly protein AP180, AP180-CT. The construct contains the clathrin-binding domain at C-terminal region of AP180, and its expression is known to inhibit clathrin-mediated endocytosis [73, 74]. We transiently transfected N2A cells with AP180 full-length wild type or a dominant-negative mutant AP180-CT. Transiently transfected N2A cells expressed AP180 and AP180-CT as detected by Western blot analysis (Figure 1B inset). However, the AP180-CT mutant did not inhibit cell toxicity induced by oA $\beta$ 42, similar to the wild type control (Figure 1B). Furthermore, in both AP180 and AP180-CT mutant transfected cells, there were similar levels of intracellular A $\beta$  accumulation as detected by immunofluorescence quantitation (figure 2B). Third, chlorpromazine, a cationic amphiphilic drug that inhibits the formation of clathrin-coated pits [75], was tested. While this compound is

toxic to N2A cells at high concentration, treatment with 2mM chlorpromazine retained 90% cell viability. Again, this treatment failed to block oA $\beta$ 42 neurotoxicity (data not shown). These combined results strongly suggest that under our experimental conditions, the clathrin-dependent endocytic pathway does not participate in oA $\beta$ 42-induced neurotoxicity.

### **Dynamin mediates oA $\beta$ 42 neurotoxicity and intracellular accumulation**

Dynamin mediates both clathrin-dependent and -independent endocytosis (for review, [76]) (refer to Figure 4). We first used two pharmacological inhibitors to block the dynamin-dependent endocytic pathway, genistein (a general tyrosine kinase inhibitor) and PP2 (a Src family tyrosine kinase inhibitor). As shown in Figure 2A, genistein significantly inhibited oA $\beta$ 42-induced toxicity. The result was consistent with previous reports that this agent protected from A $\beta$  induced toxicity in cultured hippocampal neurons [77] and SH-SY5Y cells [78]. PP2 pre-treatment also decreased A $\beta$ 42 toxicity though to a lesser extent than genistein (data not shown). In addition, we used a dominant-negative dynamin mutant K44A, an established reagent to specifically abolish dynamin function [79]. Both dynamin K44A mutant and wild type proteins were expressed in the transfected cells, as determined by Western blot analysis (Figure 2B inset). With oA $\beta$ 42 treatment, the K44A mutant inhibited neurotoxicity compared to wild type dynamin (Figure 2B). Thus, the prediction would be that blocking dynamin mediated oA $\beta$ 42 endocytosis would decrease intracellular A $\beta$  accumulation. N2A cells were transfected with dynamin wild type or mutant K44A. Treatment of these transfected cells with oA $\beta$ 42 resulted in significantly less intracellular A $\beta$  in dynamin mutant cells compared to dynamin wild type cells (Figure 2C). These data collectively support a role for dynamin in oA $\beta$ 42 endocytosis and neurotoxicity.

### **RhoA regulates oA $\beta$ 42 neurotoxicity and intracellular accumulation**

We next determined the role of RhoA in oA $\beta$ 42 neurotoxicity and endocytosis (refer to Figure 4). The small GTPase RhoA regulates the clathrin-independent endocytic pathways [76]. For these experiments, we transiently transfected N2A cells with vector alone, RhoA wild type or a dominant-negative RhoA mutant T19N [80].

Western blot analysis confirmed expression of RhoA and T19N proteins (Figure 3A inset). In cells treated with oA $\beta$ 42, the T19N RhoA mutant significantly protected cells from oA $\beta$ 42-induced neurotoxicity compare to RhoA wild type (Figure 3A). To confirm that RhoA is involved in oA $\beta$ 42 endocytosis, we assessed accumulation of intracellular A $\beta$  by treating these transfected cells oA $\beta$ 42. Significantly less A $\beta$  was detected in RhoA mutant positive cells compared to RhoA wild type cells (Figure 3B). These results suggest that RhoA is involved in oA $\beta$ 42 endocytosis and neurotoxicity.

Taken together, these data strongly suggest that endocytosis is critical for oA $\beta$ 42-induced neurotoxicity. This process is dependent on dynamin, but not clathrin, and further regulated by RhoA (Figure 4).

## **Discussion**

In recent years, it has become increasingly clear that soluble oA $\beta$  plays an essential role in the neuronal loss characteristic of AD pathology. Soluble oA $\beta$  could mediate neuronal dysfunction extracellularly by binding to cell surface receptors and disturbing downstream signaling pathways, leading to disruption of LTP and LTD, and eventual neuronal death. Several receptors have been reported to bind soluble A $\beta$ , including  $\alpha$ 7nAChR [81, 82], NMDAR [31] and cellular prion protein [27].

Alternatively, soluble oA $\beta$  toxicity could arise from intraneuronal accumulation as a result of impaired exocytosis or failed clearance following endocytosis. The importance of endocytosis in AD is underscored by a recent report identifying genetic variances in phosphatidylinositol-binding clathrin assembly protein (PICALM) associated with late onset AD [83]. PICALM facilitates endocytosis in hippocampal neurons and thus could play a role in A $\beta$  clearance in the brain [84]. However, the mechanisms underlying binding and subsequent signalling pathways or endocytosis leading to A $\beta$  intracellular accumulation remain poorly understood.

Although a major endocytic pathway in neurons is clathrin-dependent [85], we show by three complementary approaches that inhibition of this pathway did not inhibit oA $\beta$ 42 neurotoxicity (Figure 1). A reduced level of AP180 has been reported in AD patient brains [86]. Our data that AP180 did not mediate A $\beta$  toxicity suggests that AP180 could potentially regulate trafficking of proteins/enzymes involved in A $\beta$  production [87].

There is increasing evidence of clathrin- and caveolin-independent pathways mediating ligand-induced endocytosis [76, 88]. The large GTPase dynamin is involved in both clathrin-dependent and -independent pathways [76]. Our results suggest an important role for dynamin in oA $\beta$ 42-induced neurotoxicity and intraneuronal A $\beta$  accumulation. Interestingly, clathrin-independent but dynamin-dependent endocytosis was required for A $\beta$  internalization in sympathetic neurons *in vitro* [89]. The small monomeric GTPase RhoA regulates other clathrin-independent pathways, such as IL2-receptor endocytosis, [90]. Our data show RhoA regulates oA $\beta$ 42 endocytosis and neurotoxicity. The role of RhoA in oA $\beta$ 42-induced neurotoxicity is further supported by recent reports of potential roles for this GTPase in AD. For example, RhoA had an altered subcellular localization in both AD and

APP transgenic Tg2576 mouse brains [91]. Further, RhoA levels increased specifically around amyloid plaques in these models [92].

As neurons do not express caveolin-1, the principal structural protein in caveolae, and do not have caveolae structure [93], we did not pursue this pathway. Another possible route for A $\beta$  uptake is pinocytosis. A $\beta$ 40 directly conjugated with fluorescein was taken up by neurons via diffusion in a non-saturable, energy-independent process [94]. In our experiments, ATP levels were used as a measurement of neurotoxicity, precluding results based on energy independence. More importantly, while we have been able to consistently label oligomers with Alexa-488 after formation and maintain conformational stability, we are unable to prepare consistent oligomeric conformations using pre-labeled fluorescein-A $\beta$ 42 [95]. In addition, comparison between A $\beta$ 40 and 42 is problematic.

Our data (summarized in the schematic shown in Figure 4) show RhoA and dynamin-dependent steps involved in oA $\beta$ 42 neurotoxicity and intracellular A $\beta$  accumulation.

## **Conclusions**

Our experiments identify the initial steps of endocytosis required for oA $\beta$ 42-induced neurotoxicity and intracellular A $\beta$  accumulation. Specifically, A $\beta$ -induced neurotoxicity is dynamin-dependent and RhoA-regulated, but clathrin-independent. Further studies will be needed to identify potential steps in the endocytic pathways as therapeutic targets in AD.

## **Methods**

### **Materials**

Recombinant A $\beta$ 42 was purchased from rPeptide (Bogart, GA).

Hexafluoroisopropanol (HFIP) and anhydrous dimethyl sulfoxide (DMSO) were purchased from Sigma-Aldrich. Phenol-red free Ham's F12 media was obtained from Promocell (Heidelberg, Germany) and supplemented with L-glutamine (146mg/L) prior to use. Genistein and chlorpromazine were purchased from Sigma. PP2 was purchased from EMD Biosciences.

### **Oligomer formation conditions**

Oligomer preparations of A $\beta$ 42 were formed according to our previously established protocols [18, 19]. Briefly, following evaporation of HFIP in a fume hood overnight, the resulting A $\beta$ 42 peptide film was stored desiccated at -20°C. Immediately prior to use, the films were allowed to come to room temperature, solubilized to 5mM in anhydrous DMSO, sonicated in a bath sonicator (Branson) for 10 minutes, diluted to 100 $\mu$ M in phenol-red free Hams F12, and stored at 4°C for 24 hours. Oligomeric A $\beta$ 42 morphology was routinely confirmed by atomic force microscope [19].

### **Cell culture and cell viability assay**

Mouse neuroblastoma, N2A cells (ATCC) were maintained in MEM (ATCC) supplemented with 10% FBS, 2mM L-glutamine, 100U/ml of penicillin, 100ug/ml of streptomycin, as previously described [18, 69]. 5000 cells per well were plated on to 96-well plates 24 hours prior to treatment to allow attachment. Cells were then treated 10mM oA $\beta$ 42 in DMEM medium without phenol red and with 1% N2 supplement (Invitrogen). At the end of the experiment (24 hours post-treatment), cell viability was

assessed by relative cellular ATP levels using CellTiter-Glo assay kit (Promega) according to the manufacture's instruction. Statistical significance was established at  $p < 0.01$  by One-way ANOVA with Tukey test for comparison in different groups.

### **Pharmacological inhibition of endocytosis**

N2A cells were treated with pharmacological inhibitors that block specific steps during endocytosis. Pilot experiments were performed to find inhibitor concentrations that did not significantly compromise cell viability as inhibitors for cellular endocytosis could adversely affect cell viability. For example, as has been reported, chlorpromazine at higher concentration ( $10^{-4}$ - $10^{-3}$  M) killed cells, while at lower concentration ( $10^{-6}$ - $10^{-5}$  M) inhibited  $Ca^{++}$ -mediated toxicity in a neuroblastoma cell line [96]. Genistein substantially inhibited the growth of N2A cells in a dose-dependent manner with an IC50 value of 18mM, and PP2 at 3mM was lethal to the N2A cells [97]. Chlorpromazine (2mM), genistein (10mM), PP2 (1mM) were added to cell cultures at indicated concentrations 1 hr before  $\alpha A\beta 42$  treatment in DMEM with 1% N2 supplement. The final concentration of vehicle (DMSO) was 0.05% in all cultures.

### **Genetic manipulation of selected endocytic pathway proteins**

To block specific routes in the endocytic pathways, we blocked the function of key proteins in the endocytic pathway by either expressing dominant-negative proteins, or knock-down of endogenous proteins. N2A cells were transiently transfected with expression plasmids or siRNA using LipofectAmine 2000 (Invitrogen). The following endocytic proteins were transfected for expression: Rat wild type (WT) Dynamin and the dominant-negative dynamin mutant K44A (myc-tag), the dominant-negative AP180 mutant AP180-CT (Flag-tag), and RhoA WT and dominant-negative

mutant T19N (HA-tag). These plasmids were kindly provided by Dr. R. Minshall (UIC, dynamin) and Dr. L. Greene (NIH, AP180), or purchased from Missouri S&T (RhoA). To achieve highest possible transfection efficiency, we tested several transfection reagents (such as LipofectAmine and PLUS reagent, GenJet, and LipofectAmine 2000) and transfection conditions (cell density, pH of the medium and transfection incubation duration) with EGFP expression plasmid. We obtained the highest transfection efficiency with LipofectAmine 2000 at cell density of 90-100% confluence.

Small interfering RNA (siRNA) for the clathrin heavy chain (CLTC, SMARTpool L-004001-00-0005) and control Non-Targeting siRNA were purchased from Dharmacon. Cells were transfected at 20pmol siRNA in 24-well culture plates using LipofectAmine 2000 according to vendor's recommended transfection protocol. A second transfection was done the next day. These transfected cells were then split and seeded on to 96-well plate in MEM with 10% FBS. 48 hours post-transfection, these cells were treated with  $\alpha$ A $\beta$ 42 for 24 hours.

### **SDS-PAGE/Western blot characterization of targeted endocytic proteins**

Transfected cells treated in parallel to those used for cell viability assays were lysed by 15-minute incubation in RIPA buffer (50mM Tris-HCl, pH 8.0, with 150mM sodium chloride, 1.0% Igepal CA-630 (NP-40), 0.5% sodium deoxycholate, and 0.1% SDS, Sigma-Aldrich) containing protease inhibitors (Protease Inhibitor Cocktail Set I, Calbiochem), followed by centrifugation. Equal amounts of total protein were analyzed for levels of indicated proteins by Western blot analysis following SDS-PAGE using 4-12% Bis-Tris 1.5mm NuPAGE precast gels (Invitrogen). Supernatants were mixed with LDS sample buffer (Invitrogen) and electrophoresed at 90-100V for 80-90 minutes. Proteins were transferred to 0.2 $\mu$ m polyvinylidene difluoride

membranes. Membranes were blocked for 1 hour in a solution of 5% nonfat dry milk in Tris-buffered saline containing 0.0625% Tween-20 prior to incubation with primary antibody solutions. Molecular mass was estimated using pre-stained molecular weight markers (Invitrogen). CLTC was detected using a mouse anti-CLTC monoclonal antibody (C1860, Sigma; 1:1,000), myc tagged dynamin with anti-myc antibody 9E10 (Sigma; 1:5000), Flag-tagged AP180-CT with Flag antibody M2 (Sigma, 1:5000), HA tagged RhoA with rat HA antibody 3F10 (Roche Applied Science; 1:2500), and appropriate horse radish peroxidase conjugated secondary antibody. Actin, as detected with rabbit anti-actin antibody (Sigma; 1:5000), was used as total lysate loading control. Proteins were visualized with enhanced chemiluminescence Western blotting substrate (Pierce) on the Kodak 4000R imaging system.

#### **Cellular uptake of oligomeric A $\beta$ 42**

Intracellular A $\beta$  was detected by immunofluorescence analysis using a rabbit polyclonal anti-A $\beta$ 42 specific antibody (Invitrogen). N2A cells or transiently transfected N2A cells (24 hours post transfection) were seeded at 20,000 cells/well on poly-D-lysine glass coverslips in phenol-red free DMEM + 10% FBS overnight. Recombinant oA $\beta$ 42 was added to cells in DMEM medium and incubated for 30min at 37°C. At the end of the treatment, cells were washed with PBS. Cell surface bound oA $\beta$  was striped off in a solution of 0.2M acetic acid and 0.5M NaCl. Cells were fixed in 4% paraformaldehyde for 20 minutes at room temperature. Cells were permeablized with 0.3% Triton X-100 in 1xPBS for 5 minutes, and blocked for 15 minutes with 3% BSA, incubated overnight with rabbit anti-A $\beta$ 42 (1:100) at 4°C, followed by 1 hour incubation at room temperature with Alexa488-labeled donkey-anti-rabbit IgG (1:500, Invitrogen). Transfected cells were identified by co-staining

with anti-Flag antibody (M2, 1:250, Sigma) for AP180, rat anti-HA (3F10, Roche) for RhoA, or mouse anti-myc antibody 9E10 (1:200, Abcam) for dynamin, and appropriate 2<sup>nd</sup> antibody conjugated with Alexa594 (all from Invitrogen). Coverslips were mounted with Prolong Gold antifade reagent with DAPI (Invitrogen) fluorescence mounting medium on glass slides. Confocal laser scanning microscopy images were acquired on a Zeiss LSM 510 META, Axiovert 200M laser scanning confocal microscope using a Plan-Apochromate Zeiss 40X/1.3 oil immersion objective. Mean brightness of A $\beta$  signals were quantified with NIH image software.

## **Competing interests**

The authors declare that they have no competing interests.

## **Authors' contributions**

CY designed and performed the experiments, collected and analyzed the data, and wrote the manuscript. EN and KL collected and analyzed the data. MJL contributed to experimental design and data analyses, and preparation of the manuscript. All authors read and approved the final manuscript.

## **Acknowledgements**

We thank Lisa Jungbauer, Katie Younmans, and Jennifer Graham for technique assistance and manuscript preparation. This work was supported by grants from the Alzheimer's Association NIRG-06-26957 (CY); and ZEN-08-899000 and NIH/NIA

PO1AG021184 (MJL). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “*advertisement*” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

## References

1. Hardy JA, Higgins GA: Alzheimer's disease: the amyloid cascade hypothesis. *Science* 1992, 256(5054):184-185.
2. Kuo YM, Emmerling MR, Vigo-Pelfrey C, Kasunic TC, Kirkpatrick JB, Murdoch GH, Ball MJ, Roher AE: Water-soluble Abeta (N-40, N-42) oligomers in normal and Alzheimer disease brains. *J Biol Chem* 1996, 271(8):4077-4081.
3. McLean CA, Cherny RA, Fraser FW, Fuller SJ, Smith MJ, Beyreuther K, Bush AI, Masters CL: Soluble pool of Abeta amyloid as a determinant of severity of neurodegeneration in Alzheimer's disease. *Ann Neurol* 1999, 46(6):860-866.
4. Roher AE, Chaney MO, Kuo YM, Webster SD, Stine WB, Haverkamp LJ, Woods AS, Cotter RJ, Tuohy JM, Krafft GA *et al*: Morphology and toxicity of Abeta-(1-42) dimer derived from neuritic and vascular amyloid deposits of Alzheimer's disease. *J Biol Chem* 1996, 271(34):20631-20635.
5. Funato H, Enya M, Yoshimura M, Morishima-Kawashima M, Ihara Y: Presence of sodium dodecyl sulfate-stable amyloid beta-protein dimers in the hippocampus CA1 not exhibiting neurofibrillary tangle formation. *Am J Pathol* 1999, 155(1):23-28.
6. Lesne S, Koh MT, Kotilinek L, Kaye R, Glabe CG, Yang A, Gallagher M, Ashe KH: A specific amyloid-beta protein assembly in the brain impairs memory. *Nature* 2006, 440(7082):352-357.
7. Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, Rowan MJ, Selkoe DJ: Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature* 2002, 416(6880):535-539.
8. Lue LF, Kuo YM, Roher AE, Brachova L, Shen Y, Sue L, Beach T, Kurth JH, Rydel RE, Rogers J: Soluble amyloid beta peptide concentration as a predictor of synaptic change in Alzheimer's disease. *American Journal Of Pathology* 1999, 155(3):853-862.
9. Mucke L, Masliah E, Yu GQ, Mallory M, Rockenstein EM, Tatsuno G, Hu K, Kholodenko D, Johnson-Wood K, McConlogue L: High-level neuronal expression of abeta 1-42 in wild-type human amyloid protein precursor transgenic mice: synaptotoxicity without plaque formation. *J Neurosci* 2000, 20(11):4050-4058.
10. Naslund J, Haroutunian V, Mohs R, Davis KL, Davies P, Greengard P, Buxbaum JD: Correlation between elevated levels of amyloid beta-peptide in the brain and cognitive decline. *Jama* 2000, 283(12):1571-1577.

11. Spires TL, Meyer-Luehmann M, Stern EA, McLean PJ, Skoch J, Nguyen PT, Bacskai BJ, Hyman BT: Dendritic spine abnormalities in amyloid precursor protein transgenic mice demonstrated by gene transfer and intravital multiphoton microscopy. *J Neurosci* 2005, 25(31):7278-7287.
12. Moolman DL, Vitolo OV, Vonsattel JP, Shelanski ML: Dendrite and dendritic spine alterations in Alzheimer models. *J Neurocytol* 2004, 33(3):377-387.
13. Walsh DM, Tseng BP, Rydel RE, Podlisny MB, Selkoe DJ: The oligomerization of amyloid beta-protein begins intracellularly in cells derived from human brain. *Biochemistry* 2000, 39(35):10831-10839.
14. Podlisny MB, Walsh DM, Amarante P, Ostaszewski BL, Stimson ER, Maggio JE, Teplow DB, Selkoe DJ: Oligomerization of endogenous and synthetic amyloid beta-protein at nanomolar levels in cell culture and stabilization of monomer by Congo red. *Biochemistry* 1998, 37(11):3602-3611.
15. Harper JD, Wong SS, Lieber CM, Lansbury PT: Observation of metastable Abeta amyloid protofibrils by atomic force microscopy. *Chem Biol* 1997, 4(2):119-125.
16. Hartley DM, Walsh DM, Ye CP, Diehl T, Vasquez S, Vassilev PM, Teplow DB, Selkoe DJ: Protofibrillar intermediates of amyloid beta-protein induce acute electrophysiological changes and progressive neurotoxicity in cortical neurons. *J Neurosci* 1999, 19(20):8876-8884.
17. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Rozovsky I, Trommer B, Viola KL *et al*: Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A* 1998, 95(11):6448-6453.
18. Dahlgren KN, Manelli AM, Stine WB, Jr., Baker LK, Krafft GA, LaDu MJ: Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability. *J Biol Chem* 2002, 277(35):32046-32053.
19. Stine WB, Jr., Dahlgren KN, Krafft GK, LaDu MJ: In vitro characterization of conditions for amyloid-beta peptide oligomerization and fibrillogenesis. *J Biol Chem* 2003, 278:11612-11622.
20. Kaye R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, Glabe CG: Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science* 2003, 300(5618):486-489.
21. Kaye R, Head E, Sarsoza F, Saing T, Cotman CW, Necula M, Margol L, Wu J, Breydo L, Thompson JL *et al*: Fibril specific, conformation dependent antibodies recognize a generic epitope common to amyloid fibrils and fibrillar oligomers that is absent in prefibrillar oligomers. *Mol Neurodegener* 2007, 2:18.
22. Walsh DM, Lomakin A, Benedek GB, Condron MM, Teplow DB: Amyloid beta-protein fibrillogenesis. Detection of a protofibrillar intermediate. *J Biol Chem* 1997, 272(35):22364-22372.

23. Barghorn S, Nimmrich V, Striebinger A, Krantz C, Keller P, Janson B, Bahr M, Schmidt M, Bitner RS, Harlan J *et al*: Globular amyloid beta-peptide oligomer - a homogenous and stable neuropathological protein in Alzheimer's disease. *Journal of neurochemistry* 2005, 95(3):834-847.
24. Ono K, Condrón MM, Teplow DB: Structure-neurotoxicity relationships of amyloid beta-protein oligomers. *Proceedings of the National Academy of Sciences of the United States of America* 2009, 106(35):14745-14750.
25. Haass C, Selkoe DJ: Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid beta-peptide. *Nat Rev Mol Cell Biol* 2007, 8(2):101-112.
26. Wang HY, Stucky A, Liu J, Shen C, Trocme-Thibierge C, Morain P: Dissociating beta-amyloid from alpha 7 nicotinic acetylcholine receptor by a novel therapeutic agent, S 24795, normalizes alpha 7 nicotinic acetylcholine and NMDA receptor function in Alzheimer's disease brain. *J Neurosci* 2009, 29(35):10961-10973.
27. Lauren J, Gimbel DA, Nygaard HB, Gilbert JW, Strittmatter SM: Cellular prion protein mediates impairment of synaptic plasticity by amyloid-beta oligomers. *Nature* 2009, 457(7233):1128-1132.
28. De Felice FG, Velasco PT, Lambert MP, Viola K, Fernandez SJ, Ferreira ST, Klein WL: Abeta oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate receptor-dependent mechanism that is blocked by the Alzheimer drug memantine. *The Journal of biological chemistry* 2007, 282(15):11590-11601.
29. Deshpande A, Kawai H, Metherate R, Glabe CG, Busciglio J: A role for synaptic zinc in activity-dependent Abeta oligomer formation and accumulation at excitatory synapses. *The Journal of neuroscience* 2009, 29(13):4004-4015.
30. Almeida CG, Tampellini D, Takahashi RH, Greengard P, Lin MT, Snyder EM, Gouras GK: Beta-amyloid accumulation in APP mutant neurons reduces PSD-95 and GluR1 in synapses. *Neurobiol Dis* 2005, 20(2):187-198.
31. Snyder EM, Nong Y, Almeida CG, Paul S, Moran T, Choi EY, Nairn AC, Salter MW, Lombroso PJ, Gouras GK *et al*: Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci* 2005, 8(8):1051-1058.
32. Chang EH, Savage MJ, Flood DG, Thomas JM, Levy RB, Mahadomrongkul V, Shirao T, Aoki C, Huerta PT: AMPA receptor downscaling at the onset of Alzheimer's disease pathology in double knockin mice. *Proceedings of the National Academy of Sciences of the United States of America* 2006, 103(9):3410-3415.
33. Hsieh H, Boehm J, Sato C, Iwatsubo T, Tomita T, Sisodia S, Malinow R: AMPAR removal underlies Abeta-induced synaptic depression and dendritic spine loss. *Neuron* 2006, 52(5):831-843.

34. Gamblin TC, Chen F, Zambrano A, Abraha A, Lagalwar S, Guillozet AL, Lu M, Fu Y, Garcia-Sierra F, LaPointe N *et al*: Caspase cleavage of tau: Linking amyloid and neurofibrillary tangles in Alzheimer's disease. *Proc Natl Acad Sci U S A* 2003.
35. D'Andrea MR, Reiser PA, Polkovitch DA, Gumula NA, Branchide B, Hertzog BM, Schmidheiser D, Belkowski S, Gastard MC, Andrade-Gordon P: The use of formic acid to embellish amyloid plaque detection in Alzheimer's disease tissues misguides key observations. *Neurosci Lett* 2003, 342(1-2):114-118.
36. D'Andrea MR, Nagele RG, Wang HY, Lee DH: Consistent immunohistochemical detection of intracellular beta-amyloid42 in pyramidal neurons of Alzheimer's disease entorhinal cortex. *Neurosci Lett* 2002, 333(3):163-166.
37. Grundke-Iqbal I, Iqbal K, George L, Tung YC, Kim KS, Wisniewski HM: Amyloid protein and neurofibrillary tangles coexist in the same neuron in Alzheimer disease. *Proc Natl Acad Sci U S A* 1989, 86(8):2853-2857.
38. LaFerla FM, Troncoso JC, Strickland DK, Kawas CH, Jay G: Neuronal cell death in Alzheimer's disease correlates with apoE uptake and intracellular Abeta stabilization. *J Clin Invest* 1997, 100(2):310-320.
39. Takahashi RH, Milner TA, Li F, Nam EE, Edgar MA, Yamaguchi H, Beal MF, Xu H, Greengard P, Gouras GK: Intraneuronal Alzheimer abeta42 accumulates in multivesicular bodies and is associated with synaptic pathology. *Am J Pathol* 2002, 161(5):1869-1879.
40. Mori C, Spooner ET, Wisniewsk KE, Wisniewski TM, Yamaguch H, Saido TC, Tolan DR, Selkoe DJ, Lemere CA: Intraneuronal Abeta42 accumulation in Down syndrome brain. *Amyloid* 2002, 9(2):88-102.
41. Gyure KA, Durham R, Stewart WF, Smialek JE, Troncoso JC: Intraneuronal abeta-amyloid precedes development of amyloid plaques in Down syndrome. *Arch Pathol Lab Med* 2001, 125(4):489-492.
42. Busciglio J, Pelsman A, Wong C, Pigino G, Yuan M, Mori H, Yankner BA: Altered metabolism of the amyloid beta precursor protein is associated with mitochondrial dysfunction in Down's syndrome. *Neuron* 2002, 33(5):677-688.
43. Gouras GK, Tsai J, Naslund J, Vincent B, Edgar M, Checler F, Greenfield JP, Haroutunian V, Buxbaum JD, Xu H *et al*: Intraneuronal Abeta42 accumulation in human brain. *Am J Pathol* 2000, 156(1):15-20.
44. Langui D, Girardot N, El Hachimi KH, Allinquant B, Blanchard V, Pradier L, Duyckaerts C: Subcellular topography of neuronal Abeta peptide in APPxPS1 transgenic mice. *Am J Pathol* 2004, 165(5):1465-1477.
45. D'Andrea MR, Nagele RG, Wang HY, Peterson PA, Lee DH: Evidence that neurones accumulating amyloid can undergo lysis to form amyloid plaques in Alzheimer's disease. *Histopathology* 2001, 38(2):120-134.

46. Wirths O, Multhaup G, Czech C, Blanchard V, Moussaoui S, Tremp G, Pradier L, Beyreuther K, Bayer TA: Intraneuronal Abeta accumulation precedes plaque formation in beta-amyloid precursor protein and presenilin-1 double-transgenic mice. *Neurosci Lett* 2001, 306(1-2):116-120.
47. Oddo S, Caccamo A, Shepherd JD, Murphy MP, Golde TE, Kaye R, Metherate R, Mattson MP, Akbari Y, LaFerla FM: Triple-transgenic model of Alzheimer's disease with plaques and tangles: intracellular Abeta and synaptic dysfunction. *Neuron* 2003, 39(3):409-421.
48. Shie FS, LeBoeuf RC, Jin LW: Early intraneuronal Abeta deposition in the hippocampus of APP transgenic mice. *Neuroreport* 2003, 14(1):123-129.
49. Takahashi RH, Almeida CG, Kearney PF, Yu F, Lin MT, Milner TA, Gouras GK: Oligomerization of Alzheimer's beta-amyloid within processes and synapses of cultured neurons and brain. *J Neurosci* 2004, 24(14):3592-3599.
50. Skovronsky DM, Doms RW, Lee VM: Detection of a novel intraneuronal pool of insoluble amyloid beta protein that accumulates with time in culture. *J Cell Biol* 1998, 141(4):1031-1039.
51. Mazur-Kolecka B, Frackowiak J, Carroll RT, Wisniewski HM: Accumulation of Alzheimer amyloid-beta peptide in cultured myocytes is enhanced by serum and reduced by cerebrospinal fluid. *J Neuropathol Exp Neurol* 1997, 56(3):263-272.
52. Casas C, Sergeant N, Itier JM, Blanchard V, Wirths O, van der Kolk N, Vingtdoux V, van de Steeg E, Ret G, Canton T *et al*: Massive CA1/2 neuronal loss with intraneuronal and N-terminal truncated Abeta42 accumulation in a novel Alzheimer transgenic model. *Am J Pathol* 2004, 165(4):1289-1300.
53. Mochizuki A, Tamaoka A, Shimohata A, Komatsuzaki Y, Shoji S: Abeta42-positive non-pyramidal neurons around amyloid plaques in Alzheimer's disease. *Lancet* 2000, 355(9197):42-43.
54. Oddo S, Caccamo A, Tran L, Lambert MP, Glabe CG, Klein WL, Laferla FM: Temporal profile of Abeta oligomerization in an in vivo model of Alzheimer's disease: A link between Abeta and tau pathology. *J Biol Chem* 2005.
55. Kawarabayashi T, Shoji M, Younkin LH, Wen-Lang L, Dickson DW, Murakami T, Matsubara E, Abe K, Ashe KH, Younkin SG: Dimeric amyloid beta protein rapidly accumulates in lipid rafts followed by apolipoprotein E and phosphorylated tau accumulation in the Tg2576 mouse model of Alzheimer's disease. *J Neurosci* 2004, 24(15):3801-3809.
56. Oshima N, Morishima-Kawashima M, Yamaguchi H, Yoshimura M, Sugihara S, Khan K, Games D, Schenk D, Ihara Y: Accumulation of amyloid beta-protein in the low-density membrane domain accurately reflects the extent of beta-amyloid deposition in the brain. *Am J Pathol* 2001, 158(6):2209-2218.
57. Moechars D, Dewachter I, Lorent K, Reverse D, Baekelandt V, Naidu A, Tesseur I, Spittaels K, Haute CV, Checler F *et al*: Early phenotypic changes in

- transgenic mice that overexpress different mutants of amyloid precursor protein in brain. *J Biol Chem* 1999, 274(10):6483-6492.
58. Hsia AY, Masliah E, McConlogue L, Yu GQ, Tatsuno G, Hu K, Kholodenko D, Malenka RC, Nicoll RA, Mucke L: Plaque-independent disruption of neural circuits in Alzheimer's disease mouse models. *Proc Natl Acad Sci U S A* 1999, 96(6):3228-3233.
  59. Chapman PF, White GL, Jones MW, Cooper-Blacketer D, Marshall VJ, Irizarry M, Younkin L, Good MA, Bliss TVP, Hyman BT *et al*: Impaired synaptic plasticity and learning in aged amyloid precursor protein transgenic mice. *Nature Genet* 1999, 2:271-276.
  60. Zerbinatti CV, Wahrle SE, Kim H, Cam JA, Bales K, Paul SM, Holtzman DM, Bu G: Apolipoprotein E and low density lipoprotein receptor-related protein facilitate intraneuronal Abeta42 accumulation in amyloid model mice. *The Journal of biological chemistry* 2006, 281(47):36180-36186.
  61. Oddo S, Billings L, Kesslak JP, Cribbs DH, LaFerla FM: Abeta immunotherapy leads to clearance of early, but not late, hyperphosphorylated tau aggregates via the proteasome. *Neuron* 2004, 43(3):321-332.
  62. Billings LM, Oddo S, Green KN, McGaugh JL, Laferla FM: Intraneuronal Abeta causes the onset of early Alzheimer's disease-related cognitive deficits in transgenic mice. *Neuron* 2005, 45(5):675-688.
  63. Gimenez-Llort L, Blazquez G, Canete T, Johansson B, Oddo S, Tobena A, Laferla FM, Fernandez-Teruel A: Modeling behavioral and neuronal symptoms of Alzheimer's disease in mice: A role for intraneuronal amyloid. *Neurosci Biobehav Rev* 2006.
  64. Shie FS, LeBoeuf RC, Jin LW: Early intraneuronal Abeta deposition in the hippocampus of APP transgenic mice.[erratum appears in *Neuroreport*. 2004 Aug 26;15(12):1993 Note: LeBoeur, Renee C [corrected to LeBoeuf, Renee C]]. *Neuroreport* 2003, 14(1):123-129.
  65. Glabe C: Intracellular mechanisms of amyloid accumulation and pathogenesis in Alzheimer's disease. *J Mol Neurosci* 2001, 17(2):137-145.
  66. Zhang Y, McLaughlin R, Goodyer C, LeBlanc A: Selective cytotoxicity of intracellular amyloid beta peptide1-42 through p53 and Bax in cultured primary human neurons. *J Cell Biol* 2002, 156(3):519-529.
  67. Pigino G, Morfini G, Atagi Y, Deshpande A, Yu C, Jungbauer L, LaDu M, Busciglio J, Brady S: Disruption of fast axonal transport is a pathogenic mechanism for intraneuronal amyloid beta. *Proc Natl Acad Sci U S A* 2009, 106(14):5907-5912.
  68. Manelli AM, Bulfinch LC, Sullivan PM, LaDu MJ: Abeta42 neurotoxicity in primary co-cultures: effect of apoE isoform and Abeta conformation. *Neurobiol Aging* 2007, 28(8):1139-1147.

69. Manelli AM, Stine WB, Van Eldik LJ, LaDu MJ: ApoE and Abeta1-42 interactions: effects of isoform and conformation on structure and function. *J Mol Neurosci* 2004, 23(3):235-246.
70. Trommer BL, Shah C, Yun SH, Gamkrelidze G, Pasternak ES, Stine WB, Manelli A, Sullivan P, Pasternak JF, LaDu MJ: ApoE isoform-specific effects on LTP: blockade by oligomeric amyloid-beta1-42. *Neurobiology of disease* 2005, 18(1):75-82.
71. Reed MN, Hofmeister JJ, Jungbauer L, Welzel AT, Yu C, Sherman MA, Lesne S, Ladu MJ, Walsh DM, Ashe KH *et al*: Cognitive effects of cell-derived and synthetically derived Abeta oligomers. *Neurobiology of aging* 2009.
72. White JA, Manelli AM, Holmberg KH, Van Eldik LJ, LaDu MJ: Differential effects of oligomeric and fibrillar amyloid-beta1-42 on astrocyte-mediated inflammation. *Neurobiol Dis* 2005, 18(3):459-465.
73. Zhao X, Greener T, Al-Hasani H, Cushman SW, Eisenberg E, Greene LE: Expression of auxilin or AP180 inhibits endocytosis by mislocalizing clathrin: evidence for formation of nascent pits containing AP1 or AP2 but not clathrin. *J Cell Sci* 2001, 114(Pt 2):353-365.
74. Ford MG, Pearse BM, Higgins MK, Vallis Y, Owen DJ, Gibson A, Hopkins CR, Evans PR, McMahon HT: Simultaneous binding of PtdIns(4,5)P2 and clathrin by AP180 in the nucleation of clathrin lattices on membranes. *Science (New York, NY)* 2001, 291(5506):1051-1055.
75. Wang LH, Rothberg KG, Anderson RG: Mis-assembly of clathrin lattices on endosomes reveals a regulatory switch for coated pit formation. *J Cell Biol* 1993, 123(5):1107-1117.
76. Mayor S, Pagano RE: Pathways of clathrin-independent endocytosis. *Nat Rev Mol Cell Biol* 2007, 8(8):603-612.
77. Zeng H, Chen Q, Zhao B: Genistein ameliorates beta-amyloid peptide (25-35)-induced hippocampal neuronal apoptosis. *Free Radic Biol Med* 2004, 36(2):180-188.
78. Bang OY, Hong HS, Kim DH, Kim H, Boo JH, Huh K, Mook-Jung I: Neuroprotective effect of genistein against beta amyloid-induced neurotoxicity. *Neurobiol Dis* 2004, 16(1):21-28.
79. Damke H, Baba T, Warnock DE, Schmid SL: Induction of mutant dynamin specifically blocks endocytic coated vesicle formation. *J Cell Biol* 1994, 127(4):915-934.
80. Pan ZK, Ye RD, Christiansen SC, Jagels MA, Bokoch GM, Zuraw BL: Role of the Rho GTPase in bradykinin-stimulated nuclear factor-kappaB activation and IL-1beta gene expression in cultured human epithelial cells. *Journal of immunology (Baltimore, Md)* 1998, 160(6):3038-3045.

81. Nagele RG, D'Andrea MR, Anderson WJ, Wang HY: Intracellular accumulation of beta-amyloid(1-42) in neurons is facilitated by the alpha 7 nicotinic acetylcholine receptor in Alzheimer's disease. *Neuroscience* 2002, 110(2):199-211.
82. Small DH, Maksel D, Kerr ML, Ng J, Hou X, Chu C, Mehrani H, Unabia S, Azari MF, Loiacono R *et al*: The beta-amyloid protein of Alzheimer's disease binds to membrane lipids but does not bind to the alpha7 nicotinic acetylcholine receptor. *J Neurochem* 2007, 101(6):1527-1538.
83. Harold D, Abraham R, Hollingworth P, Sims R, Gerrish A, Hamshere ML, Pahwa JS, Moskvina V, Dowzell K, Williams A *et al*: Genome-wide association study identifies variants at CLU and PICALM associated with Alzheimer's disease. *Nat Genet* 2009.
84. Bushlin I, Petralia RS, Wu F, Harel A, Mughal MR, Mattson MP, Yao PJ: Clathrin assembly protein AP180 and CALM differentially control axogenesis and dendrite outgrowth in embryonic hippocampal neurons. *J Neurosci* 2008, 28(41):10257-10271.
85. Mousavi SA, Malerod L, Berg T, Kjekken R: Clathrin-dependent endocytosis. *Biochem J* 2004, 377(Pt 1):1-16.
86. Yao PJ, Morsch R, Callahan LM, Coleman PD: Changes in synaptic expression of clathrin assembly protein AP180 in Alzheimer's disease analysed by immunohistochemistry. *Neuroscience* 1999, 94(2):389-394.
87. Small SA, Gandy S: Sorting through the cell biology of Alzheimer's disease: intracellular pathways to pathogenesis. *Neuron* 2006, 52(1):15-31.
88. Kirkham M, Parton RG: Clathrin-independent endocytosis: new insights into caveolae and non-caveolar lipid raft carriers. *Biochim Biophys Acta* 2005, 1745(3):273-286.
89. Saavedra L, Mohamed A, Ma V, Kar S, de Chaves EP: Internalization of beta-amyloid peptide by primary neurons in the absence of apolipoprotein E. *The Journal of biological chemistry* 2007, 282(49):35722-35732.
90. Lamaze C, Dujeancourt A, Baba T, Lo CG, Benmerah A, Dautry-Varsat A: Interleukin 2 receptors and detergent-resistant membrane domains define a clathrin-independent endocytic pathway. *Mol Cell* 2001, 7(3):661-671.
91. Huesa G, Baltrons MA, Gomez-Ramos P, Moran A, Garcia A, Hidalgo J, Frances S, Santpere G, Ferrer I, Galea E: Altered Distribution of RhoA in Alzheimer's Disease and AbetaPP Overexpressing Mice. *J Alzheimers Dis* 2009.
92. Petratos S, Li QX, George AJ, Hou X, Kerr ML, Unabia SE, Hatzinisiriou I, Maksel D, Aguilar MI, Small DH: The beta-amyloid protein of Alzheimer's disease increases neuronal CRMP-2 phosphorylation by a Rho-GTP mechanism. *Brain* 2008, 131(Pt 1):90-108.

93. Shyng SL, Heuser JE, Harris DA: A glycolipid-anchored prion protein is endocytosed via clathrin-coated pits. *J Cell Biol* 1994, 125(6):1239-1250.
94. Kandimalla KK, Scott OG, Fulzele S, Davidson MW, Poduslo JF: Mechanism of neuronal versus endothelial cell uptake of Alzheimer's disease amyloid beta protein. *PLoS One* 2009, 4(2):e4627.
95. Jungbauer LM, Yu, C.J, Laxton, K.J., and M.J. LaDu: Preparation of fluorescently-labeled amyloid-**β** peptide assemblies: the effect of fluorophore conjugation on structure and function. . *Journal of Molecular Recognition* 2009, In Press.
96. Abe K, Sekizawa T, Kogure K: Biphasic effects of chlorpromazine on cell viability in a neuroblastoma cell line. *Neurosci Lett* 1986, 71(3):335-339.
97. Gyllberg H, Lofgren K, Lindgren H, Bedecs K: Increased Src kinase level results in increased protein tyrosine phosphorylation in scrapie-infected neuronal cell lines. *FEBS Lett* 2006, 580(11):2603-2608.

## Figures

### Figure 1 - The clathrin mediated endocytic pathway is not involved in oligomeric A $\beta$ 42-induced neurotoxicity

**A.** N2A cells were transiently transfected with siRNA for clathrin heavy chain, treated with 10 $\mu$ M oA $\beta$ 42 for 24 hours and assayed for neurotoxicity as detected with an ATP-based luminescence cell viability assay (CellTiter-Glo, Promega); no difference with treatment. **Inset**, clathrin heavy chain levels were characterized with clathrin heavy chain antibody (Sigma) by Western blot analysis with an equal amount of lysates from cells transfected with non-targeted siRNA and CLTC siRNA. **B.** N2A cells were transiently transfected with wild type AP180-FL or dominant-negative AP180-CT construct, treated with 10 $\mu$ M oA $\beta$ 42 for 24 hours, and assayed for neurotoxicity; no difference with treatment. **Inset**, AP180-CT was detected with Flag antibody (Sigma) by Western blot analysis with an equal amount of lysate from cells

transfected with AP180-FL and AP180-CT. **C.** N2A cells were transiently transfected with AP180-FL or AP180-CT. 48 hours post-transfection, cells were treated with 10  $\mu$ M oA $\beta$ 42 for 30 minutes, fixed and stained for A $\beta$  with A $\beta$ 42 specific antibody (Invitrogen, green). AP180-FL and AP180-CT mutant transfected cells were identified by anti-Flag antibody (red). Nuclei appear blue as detected by DAPI staining. Cells were individually outlined and mean fluorescence intensity of A $\beta$  signals was quantified with NIH image software. There were similar levels of A $\beta$  accumulation in these transfected cells.

## **Figure 2 - Dynamin mediates oligomeric A $\beta$ 42-induced neurotoxicity**

**A.** N2A cells were pre-treated  $\pm$  10mM genistein for 1 hour, treated with 10 $\mu$ M oA $\beta$ 42 for 24 hours  $\pm$  genistein, and assayed for neurotoxicity as described in Figure Legend 1. Significant difference ( $p < 0.01$ ) between cells  $\pm$  genistein is indicated by an asterisk (\*). **B.** N2A cells were transiently transfected with dynamin wild type, dominant-negative K44A mutant expression plasmids, or vector control; treated with 10 $\mu$ M oA $\beta$ 42 for 24 hours and assayed for neurotoxicity. Significant difference ( $p < 0.01$ ) between cells  $\pm$  dominant-negative K44A mutant are indicated by an asterisk (\*). **Inset**, expression levels of c-myc tagged dynamin were characterized by anti-myc antibody with Western blot analysis with an equal amount of lysate from cells transfected with vector, dynamin wild type, or dynamin dominant-negative K44A mutant. **C.** N2A cells were transiently transfected with dynamin wild type or dominant-negative K44A mutant expression plasmids. 48 hours post-transfection, cells were treated with 10  $\mu$ M oA $\beta$ 42 for 30 minutes, and stained for A $\beta$  with A $\beta$ 42 specific antibody (Invitrogen, green). Transfected cells were identified by anti-myc antibody (Abcam, red). Nuclei appear blue as detected by DAPI staining. Cells were individually outlined and mean fluorescence intensity of A $\beta$  signals were quantified

with NIH image software. Significant difference in A $\beta$  levels ( $p < 0.01$ ) between cells with dynamin wild type and K44A mutant is indicated by an asterisk (\*).

### **Figure 3 - RhoA mediates oligomeric A $\beta$ 42-induced neurotoxicity**

**A.** N2A cells were transiently transfected with RhoA wild type, dominant-negative T19N mutant, or vector control; treated with oA $\beta$ 42 for 24 hours; and assayed for neurotoxicity as described in Figure Legend 1. Significant difference ( $p < 0.01$ ) between cells  $\pm$  RhoA dominant-negative T19N mutant are indicated by an asterisk (\*). **Inset**, expression levels of HA-tagged RhoA were characterized by anti-HA antibody (Roche) with Western blot analysis with an equal amount of lysate from cells transfected vector, RhoA wild type or RhoA dominant-negative T19N mutant. **B.** N2A cells were transiently transfected with RhoA wild type or dominant-negative T19N mutant expression plasmids. 48 hours post-transfection, cells were treated with 10  $\mu$ M oA $\beta$ 42 for 30 minutes, fixed and stained for A $\beta$  with A $\beta$ 42 specific antibody (Invitrogen, green). RhoA wild type or T19N mutant transfected cells were identified by anti-HA antibody (red). Nuclei appear blue as detected by DAPI staining. Cells were individually outlined and mean fluorescence intensity of A $\beta$  signals were quantified with NIH image software. Significant difference ( $p < 0.01$ ) between cells transfected with RhoA wild type and T19N mutant is indicated by an asterisk (\*).

### **Figure 4 - Model for endocytic pathways mediating oligomeric A $\beta$ 42-induced neurotoxicity**

The endocytic pathways for oA $\beta$  neurotoxicity and intracellular accumulation is clathrin independent, but dynamin dependent. The pathways are further regulated by small GTPase RhoA. This figure is adapted from Mayor and Pagano [76].

Figure 1





Figure 2



**Figure 3**






Figure 4



Adapted from Mayor & Pagano (81)