

Table 2. Mean scores for the facilitators to medication adherence among HIV infected persons receiving HAART from nine facilities in Hyderabad, Andhra Pradesh, India (N=211).

I have	Mean score (SD)	Nonadherent Mean (SD)	Adherent Mean (SD)	t score, p value
...a strong sense of self-worth	3.43 (1.01)	3.00 (1.43)	3.54 (0.85)	2.33, 0.02
...seen and felt positive effects from my HIV medication	3.22 (1.19)	2.90 (1.42)	3.31 (1.11)	1.92, 0.06
...accepted my HIV diagnosis	3.63 (0.93)	3.24 (1.39)	3.73 (0.75)	2.19, 0.03
...understood the need to adhere to my HIV medication	3.46 (1.11)	2.66 (1.62)	3.66 (0.83)	3.84, <0.01
...made use of reminder tools such as pill organizers for my medicine	2.36 (1.74)	2.19 (1.80)	2.40 (1.73)	0.69, 0.49
...an HIV medicine regimen that “fits” into my daily schedule	3.16 (1.45)	2.76 (1.71)	3.26 (1.36)	1.76, 0.08
...once daily dosing of HIV medication	0.61 (1.42)	0.55 (1.37)	0.62 (1.44)	0.29, 0.77
...motivation and readiness to deal with my HIV	3.09 (1.33)	2.79 (1.49)	3.17 (1.28)	1.66, 0.10
...strong support from family and friends	3.22 (1.37)	3.14 (1.47)	3.24 (1.35)	0.40, 0.69
...gotten to know others like me who are living with HIV	3.52 (1.10)	3.33 (2.93)	3.56 (1.04)	1.06, 0.29
...regular health check-ups	3.71 (0.87)	3.48 (1.31)	3.77 (0.71)	1.41, 0.17