TITLE PAGE:

Cardiopulmonary exercise testing on a treadmill is superior than on a recumbent

ergometer in the quantification of coronary artery disease

Dejana Popovic^a, Marco Guazzi, Djordje Jakovljevic^b, Ratko Lasica^a, Marko Banovic^a,

Miodrag Ostojic^a, Ross Arena^c

^aDivision of Cardiology, University of Belgrade, Visegradska 26, 11000 Belgrade, Serbia

^bHeart Failure Unit and Cardiopulmonary Laboratory, Cardiology, I.R.C.C.S, Policlinico San

Donato, University Hospital Milan Italy;

^cInstitute of Cellular Medicine, Medical School, Newcastle University, William Leech Building,

Framlington Place, Newcastle upon Tyne NE24HH, United Kingdom

^dDepartment of Physical Therapy, College of Applied Health Sciences, University of Illinois

Chicago, 1200W Harrison St, Chicago, IL 60607, USA

RUNNING TITLE: Cardiopulmonary exercise testing modalities in ischemia

Conflict of interests:

The authors have nothing to disclose.

Word count:

Address for correspondence:

Dejana Popovic

Division of Cardiology, University of Belgrade, Serbia

Address: Veljka Dugosevica 27g, 11000 Belgrade, Serbia

Tel:00381643709684

Fax:00381113615630

E-mail: dejana popovic @yahoo.co.uk